

TGF beta 2 (Transforming growth factor-β2), Human

| Cat.no. | Product size |
|----------|--------------|
| RA-FI501 | 10µg |
| RA-FI502 | 100µg |

Storage: Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C. The protein was lyophilized from a solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5. Please use finished within one month after protein reconstitution.

Sequence:

MALDAAYCFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAGACPYLWSSDTQHSRVLSLYNTINPEASASP

CCVSQDLEPLTILYYIGKTPKIEQLSNMIVKSCKCS with polyhistidine tag at the C-terminus

Source: Escherichia coli

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography


Endotoxin Test: <0.01 EU/µg by LAL method

Suggested application: It is recommended to reconstitute the lyophilized protein in sterile H2O

to a concentration not less than 100 $\mu\text{g}/\text{mL}$ and incubate the stock solution for at least 20 min to

ensure sufficient re-dissolved.

SDS-PAGE analysis of recombinant human TGF beta 2



The product is for research only, not for diagnostic and clinical use.