

FGF-16 (Fibroblast growth factor-16), Human

| Cat.no. | Product size |
|----------|--------------|
| RA-FH401 | 25µg |
| RA-FH402 | 100µg |

Storage: Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C. The protein was lyophilized from a solution containing 1XPBS, pH 7.4. Please use finished within one month after protein reconstitution.

Sequence:

MAEVGGVFASLDWDLHGFSSSLGNVPLADSPGFLNERLGQIEGKLQRGSPTDFAHLKGILRRRQLYCRTGFHLEIFPN GTVHGTRHDHSRFGILEFISLAVGLISIRGVDSGLYLGMNERGELYGSKKLTRECVFREQFEENWYNTYASTLYKHSDSE RQYYVALNKDGSPREGYRTKRHQKFTHFLPRPV DPSKLPSMSR DLFHYR with polyhistidine tag at the

C-terminus

Source: Escherichia coli


Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin Test: <0.1 EU/µg by LAL method

Bioactivity (ED50): 30.29 ng/mL measured by its ability to induce 3T3 cells proliferation.

Suggested application: It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

SDS-PAGE analysis of recombinant human FGF-16



The product is for research only, not for diagnostic and clinical use.